2019 ACNM BENCHMARKING BEST PRACTICES

Best practices are named based on the data reported by the practice being one of the top three practices in that category. The purpose of naming "best practice" is to recognize practices that are performing well. Perhaps more importantly, it gives ACNM members the opportunity to know which practices to contact to discuss how to improve performance. By networking with "best practices", you can learn how to modify your practice processes to achieve better outcomes on quality indicators.

In an effort to provide a fair comparison, best practices are based on practice volume (total number of births per year): Low: ≤ 50 , Low Moderate: 50 -199, High Moderate: 200-499, High: ≥ 500 births/ year.

Note:

- For some volume groups, it is impossible to determine best practice for certain indicators. In the low-volume group there are many practices achieving 100% breastfeeding rates and 0% cesarean birth rates.
- There are mixed risk status and mixed models within EVERY category (e.g. high volume birth centers and low volume, high risk practices). By contacting best practices, you may find that there is a similar practice achieving better outcomes.

Congratulations to the 2019 ACNM Best Practices!

Labor and Birth Outcomes					
Variable	Low Volume ≤ 50 births/year (n=23)	Low Moderate Volume 50 -199 births/year (n=64)	High Moderate Volume 200-499 births/year (n=87)	High Volume ≥ 500 births/year (n=84)	
Spontaneous Vaginal Birth Rate	No Best Practice	 North Houston Birth Center BK Midwifery SeaMar Vancouver Women's Health Clinic 	 Willow Midwife Center for Birth and Wellness AZ, LLC Health Foundations Birth Center The Birth Center: Holistic Women's Health Care 	 Parkland Health and Hospital System Lehigh Valley Health Network, Cedar Crest Providence Medical Group Women's Services Everett 	
Primary Cesarean Rate	No Best Practice	 BK Midwifery North Houston Birth Center SeaMar Vancouver Women's Health Clinic 	 Willow Midwife Center for Birth and Wellness AZ St. Anthony Hospital in Chicago Advanced Ob-Gyn 	 Parkland Health and Hospital System HealthPartners Midwifery Services at the GW MFA 	
VBAC Success Rate	No Best Practice	No Best Practice	 Beginnings Birth Center Mayo Clinic Health System-Eau Claire Health Foundations Birth Center 	 Samaritan Obstetrics & Gynecology- Corvallis University of Iowa Midwifery Service Gundersen Healthcare Midwives 	
Rate of Cesarean Birth among Nulliparous women with a Term, Singleton baby in a Vertex position (NTSV)	No Best Practice	No Best Practice	 St. Anthony Hospital in Chicago City of Oaks Midwifery Health Foundations Birth Center 	 Intermountain Nurse-Midwives Wayne Women's Clinic CommuniCare Health Centers, Inc 	
Rate of Postpartum Hemorrhage	No Best Practice	No Best Practice	 Charlotte OB/GYN Connecticut Childbirth &Women's Center The Midwife Center for Birth & Women's Health 	 Shenandoah Women's HealthCare Women's Excellence in Midwifery Advent Health Medical Group for Women of Altamonte 	

Perineal Outcomes					
Variable	Low Volume ≤ 50 births/year (n=23) Low Moderate Volume 50 -199 births/year (n=64)		High Moderate Volume 200-499 births/year (n=87)	High Volume ≥ 500 births/year (n=84)	
Rate of women with intact perineum (perineum is intact or only small laceration(s) not requiring repair)	 Local Care Midwifery, PLLC JJB Midwifery RI Home Birth and Hope Family Health 	 Sage-Femme Midwifery Mountain Miracles Midwifery, Inc. Midwifery Care Associates 	 Willow Midwife Center for Birth and Wellness AZ Aurora UW Medical Group Midwifery & Wellness Center Mercy Women's Health 	 Midwives at Mount Auburn Hospital Wayne Women's Clinic MHealth Fairview East Certified Nurse-Midwives 	
Rate of episiotomies	No Best Practice	No Best Practice	No Best Practice	 Emory Nurse-Midwifery Practice at Grady LMG Midwifery UCSF Mission Bay 	
3rd or 4th degree laceration rate	No Best Practice	No Best Practice	 New Birth Company Overland Park LPG OBGYN Associates The Midwife Center for Birth & Women's Health 	 Intermountain Nurse-Midwives Catawba Valley Medical Center Maternity Services USF Midwifery 	

No Best Practice: Unable to establish a "Best Practice" as many practices in the birth volume category reported 0 or 100% for that variable.

Infant Outcomes					
Variable	Low Volume ≤ 50 births/year (n=23)	Low Moderate Volume 50 -199 births/year (n=64)	High Moderate Volume 200-499 births/year (n=87)	High Volume ≥ 500 births/year (n=84)	
Preterm Birth Rate (infants from a singleton birth born at less than 37weeks gestation)	No Best Practice	No Best Practice	 •Midwives at VCU Health •Health Foundations Birth Center •The Midwife Center for Birth & Women's Health 	 Kaiser Westside Medical Center Mount Sinai Midwives Atrius Health 	
Rate of Low Birth Weight (infants from a singleton birth born weighing less than 2500 grams)	No Best Practice	No Best Practice	 Silverton Health Midwives Olmsted Medical Center St. Anthony Hospital in Chicago 	 BirthCare HealthCare University of Utah College of Nursing Parkland Health and Hospital System Atrius Health 	
Rate of NICU Admissions of infants born from a singleton birth. (any admission to a level 2 or level 3 nursery for any length of time)	No Best Practice	No Best Practice	Fairhaven OB/Gyn, Inc.Harrisonburg OB/GYNLPG OBGYN Associates	Parkland Health and Hospital SystemWCCMidwives at Magee	
Exclusive Breastfeeding Rate first 48 hours No Best Practice		No Best Practice	No Best Practice	 Valley Women for Women UC San Diego Nurse-Midwifery Service Beth Israel Deaconess Plymouth OBGYN & Midwifery 	

No Best Practice: Unable to establish a "Best Practice" as many practices in the birth volume category reported 0 or 100% for that variable

Breastfeeding Outcomes				
Lo Volu Sirths Variable ≤ 5		Low Moderate Volume 50 -199 births/year (n=64)	High Moderate Volume 200-499 births/year (n=87)	High Volume ≥ 500 births/year (n=84)
Rate of Women who Attended their 6week Postpartum Visit.	No Best Practice	No Best Practice	No Best Practice	 UNC Midwives UC San Diego Nurse- Midwifery Service Lifecycle Womancare
Rate of Women Breastfeeding at their 6week Postpartum Visit. No Best Practice		No Best Practice	 Advanced Ob-Gyn Dar a Luz Birth & Health Center Willow Midwives PLLC 	 Lifecycle Womancare UC San Diego Nurse- Midwifery Service UNC Midwives

No Best Practice: Unable to establish a "Best Practice" as many practices in the birth volume category reported 0 or 100% for that variable.

Resource Utilization Outcomes				
Variable Low Volum ≤ 50 births/yo (n=23)		Low Moderate Volume 50 -199 births/year (n=64)	High Moderate Volume 200-499 births/year (n=87)	High Volume ≥ 500 births/year (n=84)
Rate of Inductions	No Best Practice	No Best Practice	 LPG OBGYN Associates Beginnings Birth Center New Birth Company Overland Park 	 Acclaim Physicians Group Parkland Health and Hospital System Women's Health Specialists and Midwives of Dayton
Rate of epidurals used for pain relief during labor (not including epidurals placed only for anesthesia for operative delivery) No Best Practice		 River and Mountain Midwives Inanna Birth & Women's Care Riverside Midwifery LLC 	 Brooklyn Birthing Center Beginnings Birth Center The Midwife Group Women's Health and Birth Center 	 Sutter Davis Women's Health Wayne Women's Clinic Women's Health at Valley View Hospital

No Best Practice: Unable to establish a "Best Practice" as many practices in the birth volume category reported 0 or 100% for that variable.